

## IMPORTANT NOTICE

Mr \_\_\_\_\_ attended the pharmacodynamic Doppler ultrasound examination today, on \_\_\_\_\_ (date). The vasoactive substance, \_\_\_\_\_  $\mu\text{g}$  of **Caverject**, was injected in the cavernous body of the penis, which can cause prolonged erection in very rare cases.

If the erection stimulated in this manner lasts for **more than 4 hours**, it is **NECESSARY** to see a doctor, a urologist if possible. If the erection lasts for more than 6 hours, it can cause lasting structural disorders of the cavernous tissue and subsequent aggravation of erectile dysfunction.

### Instructions for doctors

The initial therapy consists of the aspiration of up to 100 ml of blood from cavernous bodies in total. If this has no effect, 1 ml of adrenaline solution must be injected in the cavernous body. This procedure may be repeated several times after a few minutes' interval if the penis has not retracted within a few minutes after the injection. If the procedure is not effective, an urgent intervention of a urologist is necessary.

### Preparation of the solution:

1 ml (ampoule) of adrenaline in 9 ml of saline or 5% of glucose.

Method of injection: with the thinnest needle, perpendicular to the skin, in the central part of the left or right cavernous body. Because of systemic effect, specific care is required in patients with coronary disease, unregulated arterial hypertension and cerebral ischemia.